

<b>Product Name:</b>	Colistin Sulfate, USP
<b>Product Number:</b>	C083
<b>CAS Number:</b>	1264-72-8
<b>Molecular Formula:</b>	$C_{52}H_{98}N_{16}O_{13} \cdot 2.5 H_2SO_4$ (Colistin Sulfate A)
<b>Molecular Weight:</b>	1155.43 (Colistin Sulfate A)
<b>Form:</b>	Powder
<b>Appearance:</b>	White or almost white powder
<b>Solubility:</b>	50mg/mL H <sub>2</sub> O: Clear and Light yellow solution
<b>Source:</b>	<i>Bacillus Polymyxa subsp.</i>
<b>Elemental Analysis:</b>	As≤2 ppm
<b>Potency (on a dry basis):</b>	≥19,900 u/mg
<b>Absorbance:</b>	$A_{470} \leq 0.4$
<b>pH:</b>	4.0-7.0
<b>Optical Rotation:</b>	-63° to -73° (dried substance)
<b>Storage Conditions:</b>	2-8°C
<b>Description:</b>	<p>Colistin sulfate, USP (syn: polymyxin E) is a cationic, cyclic polypeptide antibiotic belonging to the polymyxin family. It is a natural product discovered in 1949 that is nonribosomally synthesized by <i>Bacillus polymyxa</i> subspecies <i>colistinus</i> Koyama. Colistin is mixture of Colistin A, B, and C. Colistin targets the bacterial cell membrane, leading to reduced permeability and cell death. It is effective against Gram-negative bacteria. It is soluble in water (50 mg/ml).</p> <p>TOKU-E offers three forms of Colistin:</p> <ul style="list-style-type: none"> <li>• Colistin sulfate, USP (C083)</li> <li>• <u>Colistin sulfate, EP (C039)</u></li> <li>• <u>Colistin sodium methanesulfonate (C073)</u></li> </ul> <p>This product is considered a dangerous good. Quantities above 1 g may be subject to additional shipping fees. Please contact us for details.</p>
<b>Mechanism of Action:</b>	<p>Colistin sulfate displaces essential ions (magnesium and calcium) in the lipopolysaccharide layer of the cell wall leading to local disturbance of the outer membrane, which causes increased permeability and eventually cell lysis. Colistin also has anti-endotoxin activity. The endotoxin of Gram-negative bacteria is the lipid A portion of the lipopolysaccharide molecules, and colistin binds and neutralizes the lipopolysaccharide (Falagas et al, 2005).</p>

**Spectrum:** Colistin sulfate is used primarily against Gram-negative bacteria and can also be used against Mycobacteria.

**Microbiology Applications** Colistin is commonly used in clinical *in vitro* microbiological antimicrobial susceptibility tests (panels, discs, and MIC strips) against gram negative microbial isolates. Colistin has also shown high potency against high-resistant superbug strains. Medical microbiologists use AST results to recommend antibiotic treatment options for infected patients. Representative MIC values include:

- *Shigella* spp. 64 µg/mL -128 µg/mL
- *Haemophilus influenzae* 0.4 µg/mL – 0.8 µg/mL
- For a complete list of colistin MIC values, [click here](#).

## Media Supplement

Colistin is routinely used as a selection agent in several types of isolation media:

Columbia Blood Agar - *Campylobacter* selective supplement (Butzler)

Columbia Blood Agar - *Staphylococcus/Streptococcus* selective supplement

Columbia Blood Agar - *Streptococcus* Selective Supplement

Listeria Selective Agar - *Listeria* Selective Supplement

Listeria Selective Agar - Modified *Listeria* Selective Supplement

## Plant Biology Applications

*Plasmodiophora brassicae* spores were treated with colistin sulfate and used to inoculate hairy roots of cabbage, Chinese cabbage, turnip, and rape in a useful dual culture system for *P. brassicae* (Asano et al, 1999).

## References:

Asano T, Kageyama K and Myakumachi (1999) Surface disinfection of resting spores of *Plasmodiophora brassicae* used to infect hairy roots of Brassica spp. Phytopathol. 89(4):314-319 PMID 18944777

Falagas ME and Kasiakou SK and Saravolatz LD(2005) Colistin: the revival of polymyxins for the management of multidrug-resistant gram-negative bacterial infections. Clin. Infect. Dis. 40(9):1333-1341 PMID 15825037

Koen N, van Breda SV and Loots DT (2018) Elucidating the antimicrobial mechanisms of colistin sulfate on *Mycobacterium tuberculosis* using metabolomics. Tuberculosis 111:14-19 PMID 30029899

Komura S, Kurahashi K (1979) Partial purification and properties of L-2,4-diaminobutyric acid activating enzyme from a polymyxin E producing organism. J. Biochem (Tokyo) 86:1013-1021

Leifert C, Ritchie JY and Waites WM (1991) Contaminants of plant-tissue and cell cultures. World J. Microbiol. Biotechnol. 7(4):452-469 PMID 24425131

Mueller MJ, Brodschelm W, Spannagl E and Zenk MH (1993) Signaling in the elicitation process is mediated through the octadecanoid pathway leading to jasmonic acid. Proc. Natl. Acad. Sci. USA Vol. 90(16):7490-7494 PMID 11607420

Wallace, SJ et al (2012) Interaction of colistin and colistin methanesulfonate with liposomes: colloidal aspects and implications for formulation J. Pharma Sci 101 (9):3347-3359 PMID 22623044

If you need any help, contact us: [info@toku-e.com](mailto:info@toku-e.com). Find more information on: [www.toku-e.com/](http://www.toku-e.com/)