



Blasticidin S HCl Solution (10 mg/mL in 20mM HEPES) PRODUCT DATA SHEET

issue date 07/19/2019

Product Name: Blasticidin S HCl Solution (10 mg/mL in 20mM HEPES)

Product Number: B006-B007

CAS Number: 3513-03-9

Molecular Formula: $C_{17}H_{26}N_8O_5 \cdot HCl$

Molecular Weight: 458.90 g/mol

Form: Solution (sterile)

Appearance: Clear and colorless or light yellow solution

Source: *Streptomyces griseochromogenes*

pH: 7.2-7.5

Storage Conditions: -20°C

Description: Blasticidin S HCl is a peptidyl nucleoside produced by several species of *Streptomyces* that was first isolated from *S. griseochromogenes* in 1958. Blasticidin S inhibits protein synthesis and is active against bacteria, fungi, nematodes, and tumor cells. The compound is used as a selection antibiotic for both eukaryotic and prokaryotic cells, and a marker for strain manipulation.

TOKU-E carries two forms of Blasticidin S HCl:

- Blasticidin S HCl solution (10 mg/ml in 20 mM HEPES)(B006-B007)
- Blasticidin S HCl (B001)

Blasticidin S HCl solution contains 10 mg/mL blasticidin S HCl in 20mM HEPES.

B006 (10 x 1 mL) contains 10 mg blasticidin S HCl per vial (100 mg total).

B007 (20 mL) contains 200 mg blasticidin S HCl per vial.

This product is considered a dangerous good. Quantities above 1 g may be subject to additional shipping fees.

Mechanism of Action: Blastidicin S HCl inhibits protein synthesis in prokaryotic and eukaryotic cells by binding to the ribosomal P-site which strengthens tRNA binding and slows down and prevents subsequent peptide synthesis.

Mechanisms of resistance

Resistance to Blastidicin S is conferred by *bsr*, *BSD*, and *bls* resistance genes isolated from *Bacillus cereus* K55-S1, *Aspergillus terreus*, and *Streptoverticillum* spp, respectively.

The ***bsr* resistance gene** is a 420 bp fragment and encodes a 15 kDa Blastidicin S deaminase which catalyzes the reaction of Blastidicin S to deaminohydroxyblastidicin S. Deaminohydroxyblastidicin S is a biologically inactive derivative of Blastidicin S and does not interact with or inhibit prokaryotic or eukaryotic ribosomes.

The ***bsd* resistance gene** is a 393 bp fragment and also encodes a Blastidicin S deaminase enzyme which catalyzes a similar reaction to the *BSR* deaminase. A study by Kimura et al. found the transfection frequency with *bsd* to be 80X greater than with *bsr* when using FM3A cells.

The ***bls* gene resistance gene** encodes an acetyltransferase which interacts with acetyl-coenzyme A and prevents Blastidicin S from inhibiting protein synthesis.

Microbiology Applications Blastidicin S HCl can be used as a selection agent after transformation of prokaryotic (bacterial) cells, namely *E. coli*. Optimal Blastidicin S HCl selection concentrations range from 25 - 100 µg/mL and should be tested for each experimental condition. Selective media containing Blastidicin S HCl should contain a low salt concentration (<90mM) and pH ≤7 to avoid blastidicin degradation.

References:

Adachi H, Hasebe T, Yoshinaga K , Ohta T and Sutoh K (1994) Isolation of *Dictyostelium discoideum* cytokinesis mutants by restriction enzyme-mediated integration of the Blastidicin S resistance marker. *Biochem. Biophys. Res. Comm.* 205(3):1808-1814

Bento, FM (2004) Over Expression of the Selectable Marker Blastidicin S Deaminase Gene Is Toxic to Human Keratinocytes and Murine BALB/MK Cells." *BMC Biotechnol.* 4 (29):1-10 PMID 15575952.

Izumi M. et al., 1991. Blastidicin S-resistance gene (*bsr*): A novel selectable marker for mammalian cells. *Exp.Cell Res.*197:229-33

Lu K-T et al (2016) Ovatodiolide inhibits breast cancer stem/progenitor cells through SMURF2-mediated downregulation of Hsp27. *Toxins* 8(5):127.

Kimura M, Takatsuki A, Yamaguchi I (1994) Blastidicin S deaminase gene from *Aspergillus terreus*(*BSD*): A new drug resistance gene for transfection of mammalian cells. *Biochim. Biophys. Acta.* 1219(3):653-65 PMID 7948022

Svidritskiy E, Ling C, Ermolenko DN, Korostelev AA (2013) Blastidicin S Inhibits Translation by Trapping Deformed TRNA on the Ribosome. *PNAS* 110(30):12283-12288 PMID 23824292

Takeuchi S, Hirayama K, Ueda K, Sakai H and Yonehara H (1958) Blastidicin S, a new antibiotic. *J. Antibiot.* 11(1):1-5 PMID 13525246

Yamaguchi I et al (1990) Expression of the Blastidicin S Deaminase Gene (*bsr*) in Tobacco: Fungicide Tolerance and a New Selective Marker for Transgenic Plants. *Mol. Gen. Genet* (2):332-334 PMID 2250657

If you need any help, contact us: info@toku-e.com. Find more information on: www.toku-e.com/